

Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	Books
Search	PubMed	[v] for				Go	Clear	
		[x] Limits		Preview/Index	History	Clipboard	Details	

Display	Abstract	[v]	Show	20	[v]	Sort	[v]	Send to	Text	[v]
---------	----------	-----	------	----	-----	------	-----	---------	------	-----

Entrez
PubMed

☐ 1: Plant J. 2000 Mar;21(5):445-54/

Related Articles, Links



A bifunctional epimerase-reductase acts downstream of the MUR1 gene product and completes the de novo synthesis of GDP-L-fucose in Arabidopsis.

Bonin CP, Reiter WD.

University of Connecticut, Department of Molecular and Cell Biology, Storrs, CT 06269, USA.

PubMed
Services

Related
Resources

L-Fucose is a monosaccharide found as a component of glycoproteins and cell wall polysaccharides in higher plants. The MUR1 gene of *Arabidopsis thaliana* encodes a GDP-D-mannose 4,6-dehydratase catalyzing the first step in the de novo synthesis of GDP-L-fucose from GDP-D-mannose (Bonin et al. 1997, Proc. Natl Acad. Sci. USA, 94, 2085-2090). Plant genes encoding the subsequent steps in L-fucose synthesis (3,5-epimerization and 4-reduction) have not been described previously. Based on sequence similarities to a bacterial gene involved in capsule synthesis we have cloned a gene from *Arabidopsis*, now designated GER1, which encodes a bifunctional 3, 5-epimerase-4-reductase in L-fucose synthesis. The combined action of the MUR1 and GER1 gene products converts GDP-D-mannose to GDP-L-fucose in vitro demonstrating that this entire nucleotide-sugar interconversion pathway could be reconstituted using plant genes expressed in *Escherichia coli*. In vitro assays indicated that the GER1 protein does not act as a GDP-D-mannose 3, 5-epimerase, an enzymatic activity involved in the de novo synthesis of GDP-L-galactose and L-ascorbic acid. Similarly, L-ascorbate levels in GER1 antisense plants were unchanged indicating that GDP-D-mannose 3,5-epimerase is encoded by a separate gene.

PMID: 10758496 [PubMed - indexed for MEDLINE]


Display	Abstract	[v]	Show	20	[v]	Sort	[v]	Send to	Text	[v]
---------	----------	-----	------	----	-----	------	-----	---------	------	-----

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Freedom of Information Act | Disclaimer



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	Books			
Search	PubMed	▼ for						Go	Clear		
<input checked="" type="checkbox"/> Limits Preview/Index History Clipboard Details											
Display		Abstract	▼	Show	20	▼	Sort	▼	Send to	Text	▼

Entrez
PubMed

☐ 1: J Biol Chem. 1988 Feb 5;263(4):1693-7.

Related Articles, Links

FREE full text article at
www.jbc.org

An epimerase-reductase in L-fucose synthesis.

Chang S, Duerr B, Serif G.

PubMed
Services

Department of Biochemistry, Ohio State University, Columbus 43210-1292.

Related
Resources


The first committed enzyme in GDP-L-fucose formation from GDP-D-mannose is GDP-D-mannose 4,6-dehydratase, which forms GDP-4-keto-6-deoxy-D-mannose. The uncertain enzymatic steps beyond this point were examined in this study. Assays were developed for the epimerase and reductase activities which the putative pathway would predict. A protein was isolated exhibiting homogeneity by several criteria. This single protein, which forms GDP-L-fucose from GDP-4-keto-6-deoxy-D-mannose and NADH, appears to possess both epimerase and reductase capabilities and may be termed GDP-4-keto-6-deoxy-D-mannose-3,5-epimerase-4-reductase. Analysis on a molecular sieve column using fast protein liquid chromatography established a molecular weight of 63,100 for the native enzyme, whereas sodium dodecyl sulfate-polyacrylamide gel electrophoresis established a subunit molecular weight of 31,500.

PMID: 3338988 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Show	20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	------	----	---	------	---	---------	------	---

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

02/11/1988 09:14:11



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	Books
Search	PubMed	for				Go	Clear	
		Limits		Preview/Index		History		Clipboard
		Details						
Display		Abstract		Show: 20		Sort		Send to Text

Entrez
PubMed

1: J Mol Biol. 2000 Oct 13;303(1):77-91.

Related Articles, Links

[Online Full-text](#)

**Probing the catalytic mechanism of
GDP-4-keto-6-deoxy-d-mannose Epimerase/Reductase by kinetic
and crystallographic characterization of site-specific mutants.**

PubMed
Services


Rosano C, Bisso A, Izzo G, Tonetti M, Sturla L, De Flora A, Bolognesi M.

Department of Physics-INFN and Advanced Biotechnology Center-IST,
University of Genova, Largo Rosanna Benzi 10, Genova, I-16132, Italy.

Related
Resources

GDP-4-keto-6-deoxy-d-mannose epimerase/reductase is a bifunctional enzyme responsible for the last step in the biosynthesis of GDP-l-fucose, the substrate of fucosyl transferases. Several cell-surface antigens, including the leukocyte Lewis system and cell-surface antigens in pathogenic bacteria, depend on the availability of GDP-l-fucose for their expression. Therefore, the enzyme is a potential target for therapy in pathological states depending on selectin-mediated cell-to-cell interactions. Previous crystallographic investigations have shown that GDP-4-keto-6-deoxy-d-mannose epimerase/reductase belongs to the short-chain dehydrogenase/reductase protein homology family. The enzyme active-site region is at the interface of an N-terminal NADPH-binding domain and a C-terminal domain, held to bind the substrate. The design, expression and functional characterization of seven site-specific mutant forms of GDP-4-keto-6-deoxy-d-mannose epimerase/reductase are reported here. In parallel, the crystal structures of the native holoenzyme and of three mutants (Ser107Ala, Tyr136Glu and Lys140Arg) have been investigated and refined at 1.45-1.60 Å resolution, based on synchrotron data (R-factors range between 12.6 % and 13.9 %). The refined protein models show that besides the active-site residues Ser107, Tyr136 and Lys140, whose mutations impair the overall enzymatic activity and may affect the coenzyme binding mode, side-chains capable of proton exchange, located around the expected substrate (GDP-4-keto-6-deoxy-d-mannose) binding pocket, are selectively required during the epimerization and reduction steps. Among these, Cys109 and His179 may play a primary role in proton exchange between the enzyme and the epimerization catalytic intermediates. Finally, the additional role of mutated active-site residues involved in substrate recognition and in enzyme stability has been analyzed. Copyright 2000 Academic Press.

PMID: 11021971 [PubMed - indexed for MEDLINE]



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	
Search	PubMed	▼ for					Preview	Go
		✓ Limits	Preview/Index	History	Clipboard	Details		

- Search History will be lost after eight hours of inactivity.
- To combine searches use # before search number, e.g., #2 AND #6.
- Search numbers may not be continuous; all searches are represented.

Entrez
PubMed

Search	Most Recent Queries	Time	Result
#4	Related Articles for PubMed (Select 11021971)	10:09:20	<u>188</u>
#2	Search Rosano C Field: Author, Limits: Publication Date from 2000 to 2000	10:05:46	<u>3</u>
#1	Search Camillo R Field: Author, Limits: Publication Date from 2000 to 2000	10:05:27	<u>1</u>


PubMed
Services

Clear History

Related
Resources

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

Nov 23 2001 10:09:20



Entrez	PubMed	Nucleotide	Protein	Genome	Structure	PMC	Journals	Books
Search	PubMed	for				Go	Clear	
<input checked="" type="checkbox"/> Limits		Preview/Index		History		Clipboard		Details
Display		Summary		Show: 20	Sort	Send to		Text
Items 1-20 of 188						Page 1		of 10 Next

Entrez
PubMed

- ☐ 1: [Rosano C, Bisso A, Izzo G, Tonetti M, Sturla L, De Flora A, Bolognesi M.](#) Related Articles, Links

Probing the catalytic mechanism of GDP-4-keto-6-deoxy-d-mannose Epimerase/Reductase by kinetic and crystallographic characterization of site-specific mutants.
J Mol Biol. 2000 Oct 13;303(1):77-91.
PMID: 11021971 [PubMed - indexed for MEDLINE]

PubMed
Services

- ☐ 2: [Rizzi M, Tonetti M, Vigevari P, Sturla L, Bisso A, Flora AD, Bordo D, Bolognesi M.](#) Related Articles, Links

GDP-4-keto-6-deoxy-D-mannose epimerase/reductase from Escherichia coli, a key enzyme in the biosynthesis of GDP-L-fucose, displays the structural characteristics of the RED protein homology superfamily.
Structure. 1998 Nov 15;6(11):1453-65.
PMID: 9817848 [PubMed - indexed for MEDLINE]

Related
Resources

- ☐ 3: [Somers WS, Stahl ML, Sullivan FX.](#) Related Articles, Links

GDP-fucose synthetase from Escherichia coli: structure of a unique member of the short-chain dehydrogenase/reductase family that catalyzes two distinct reactions at the same active site.
Structure. 1998 Dec 15;6(12):1601-12.
PMID: 9862812 [PubMed - indexed for MEDLINE]

- ☐ 4: [Tonetti M, Rizzi M, Vigevari P, Sturla L, Bisso A, De Flora A, Bolognesi M.](#) Related Articles, Links

















Preliminary crystallographic investigations of recombinant GDP-4-keto-6-deoxy-D-mannose epimerase/reductase from E. coli.
Acta Crystallogr D Biol Crystallogr. 1998 Jul 1;54 (Pt 4):684-6.
PMID: 9761875 [PubMed - indexed for MEDLINE]







- ☐ 5: [Chang S, Duerr B, Serif G.](#) Related Articles, Links

An epimerase-reductase in L-fucose synthesis.
J Biol Chem. 1988 Feb 5;263(4):1693-7.
PMID: 3338988 [PubMed - indexed for MEDLINE]

- ☐ 6: [Somoza JR, Menon S, Schmidt H, Joseph-McCarthy D, Dessen A, Stahl ML, Somers WS, Sullivan FX.](#) Related Articles, Links

Structural and kinetic analysis of Escherichia coli GDP-mannose 4,6 dehydratase provides insights into the enzyme's catalytic mechanism and regulation by GDP-fucose.
Structure Fold Des. 2000 Feb 15;8(2):123-35.
PMID: 10673432 [PubMed - indexed for MEDLINE]

-  **7:** Albermann C, Distler J, Piepersberg W. Related Articles, Links
 Preparative synthesis of GDP-beta-L-fucose by recombinant enzymes from enterobacterial sources.
Glycobiology. 2000 Sep;10(9):875-81.
PMID: 10988249 [PubMed - indexed for MEDLINE]
-  **8:** Mattila P, Rabina J, Hortling S, Helin J, Renkonen R. Related Articles, Links
 Functional expression of Escherichia coli enzymes synthesizing GDP-L-fucose from inherent GDP-D-mannose in Saccharomyces cerevisiae.
Glycobiology. 2000 Oct;10(10):1041-7.
PMID: 11030750 [PubMed - indexed for MEDLINE]
-  **9:** Menon S, Stahl M, Kumar R, Xu GY, Sullivan F. Related Articles, Links
 Stereochemical course and steady state mechanism of the reaction catalyzed by the GDP-fucose synthetase from Escherichia coli.
J Biol Chem. 1999 Sep 17;274(38):26743-50.
PMID: 10480878 [PubMed - indexed for MEDLINE]
-  **10:** Bonin CP, Reiter WD. Related Articles, Links
 A bifunctional epimerase-reductase acts downstream of the MUR1 gene product and completes the de novo synthesis of GDP-L-fucose in Arabidopsis.
Plant J. 2000 Mar;21(5):445-54.
PMID: 10758496 [PubMed - indexed for MEDLINE]
-  **11:** Yamamoto K, Katayama I, Onoda Y, Inami M, Kumagai H, Tochikura T. Related Articles, Links
 Evidence that the enzyme catalyzing the conversion of guanosine diphosphate D-mannose to a 4-keto sugar nucleotide intermediate requires nicotinamide adenine dinucleotide phosphate.
Arch Biochem Biophys. 1993 Feb 1;300(2):694-8.
PMID: 7679567 [PubMed - indexed for MEDLINE]
-  **12:** Tonetti M, Sturla L, Bisso A, Zanardi D, Benatti U, De Flora A. Related Articles, Links
 The metabolism of 6-deoxyhexoses in bacterial and animal cells.
Biochimie. 1998 Nov;80(11):923-31. Review.
PMID: 9893952 [PubMed - indexed for MEDLINE]
-  **13:** Bonin CP, Potter I, Vanzin GF, Reiter WD. Related Articles, Links
 The MUR1 gene of Arabidopsis thaliana encodes an isoform of GDP-D-mannose-4,6-dehydratase, catalyzing the first step in the de novo synthesis of GDP-L-fucose.
Proc Natl Acad Sci U S A. 1997 Mar 4;94(5):2085-90.
PMID: 9050909 [PubMed - indexed for MEDLINE]
-  **14:** Mulichak AM, Bonin CP, Reiter WD, Garavito RM. Related Articles, Links
 Structure of the MUR1 GDP-mannose 4,6-dehydratase from Arabidopsis thaliana: implications for ligand binding and specificity.
Biochemistry. 2002 Dec 31;41(52):15578-89.
PMID: 12501186 [PubMed - indexed for MEDLINE]

- ☐ 15: [Jarvinen N, Maki M, Rabina J, Roos C, Mattila P, Renkonen R.](#) Related Articles, Links
 Cloning and expression of *Helicobacter pylori* GDP-L-fucose synthesizing enzymes (GMD and GMER) in *Saccharomyces cerevisiae*.
Eur J Biochem. 2001 Dec;268(24):6458-64.
PMID: 11737200 [PubMed - indexed for MEDLINE]
- ☐ 16: [Suzuki N, Nakano Y, Yoshida Y, Nezu T, Terada Y, Yamashita Y, Koga T.](#) Related Articles, Links
 Guanosine diphosphate-4-keto-6-deoxy-d-mannose reductase in the pathway for the synthesis of GDP-6-deoxy-d-talose in *Actinobacillus actinomycetemcomitans*.
Eur J Biochem. 2002 Dec;269(23):5963-71.
PMID: 12444986 [PubMed - indexed for MEDLINE]
- ☐ 17: [Wolucka BA, Persiau G, Van Doorselaere J, Davey MW, Demol H, Vandekerckhove J, Van Montagu M, Zabeau M, Boerjan W.](#) Related Articles, Links
 Partial purification and identification of GDP-mannose 3",5"-epimerase of *Arabidopsis thaliana*, a key enzyme of the plant vitamin C pathway.
Proc Natl Acad Sci U S A. 2001 Dec 18;98(26):14843-8.
PMID: 11752432 [PubMed - indexed for MEDLINE]
- ☐ 18: [Ohyama C, Smith PL, Angata K, Fukuda MN, Lowe JB, Fukuda M.](#) Related Articles, Links
 Molecular cloning and expression of GDP-D-mannose-4,6-dehydratase, a key enzyme for fucose metabolism defective in Lec13 cells.
J Biol Chem. 1998 Jun 5;273(23):14582-7.
PMID: 9603974 [PubMed - indexed for MEDLINE]
- ☐ 19: [Kneidinger B, Graninger M, Adam G, Puchberger M, Kosma P, Zayni S, Messner P.](#) Related Articles, Links
 Identification of two GDP-6-deoxy-D-lyxo-4-hexulose reductases synthesizing GDP-D-rhamnose in *Aneurinibacillus thermoaerophilus* L420-91T.
J Biol Chem. 2001 Feb 23;276(8):5577-83. Epub 2000 Nov 28.
PMID: 11096116 [PubMed - indexed for MEDLINE]
- ☐ 20: [Tonetti M, Sturla L, Bisso A, Benatti U, De Flora A.](#) Related Articles, Links
 Synthesis of GDP-L-fucose by the human FX protein.
J Biol Chem. 1996 Nov 1;271(44):27274-9.
PMID: 8910301 [PubMed - indexed for MEDLINE]

Display	Summary	Show	20	Sort	Send to	Text
Items 1 - 20 of 188					Page	1 of 10 Next

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer